arXiv:math/0508086v1 [math.CO] 4 Aug 2005

ON ABELIAN $(2^{2m+1}(2^{m-1}+1), 2^m(2^m+1), 2^m)$-DIFFERENCE SETS

K. T. ARASU, YU QING CHEN AND ALEXANDER POTT

Abstract. In this paper we prove that an abelian group contains $(2^{2m+1}(2^{m-1}+1), 2^m(2^m+1), 2^m)$-difference sets with $m \geqslant 3$ if and only if it contains an elementary abelian 2-group of order 2^{2m}. Our proof shows that the method of constructing such difference sets is essentially unique.

1. Introduction

Given a finite group G of order v, a k-subset D of G is called a (v,k,λ)-difference set if for every $g \neq 1$ in G, there are exactly λ pairs of $(d_1,d_2) \in D \times D$ such that $d_1 d_2^{-1} = g$. A (v,k,λ)-difference set in a group G gives rise to a (v,k,λ)-symmetric design whose automorphism group contains G as a subgroup which acts regularly and transitively on the design. For more details on difference sets, we refer the reader to Beth et al. [3]. Much of the research on the existence or non-existence of difference sets in abelian groups has been focused on the exponent bound of the groups. The best and most complete results of this type are difference sets in abelian 2-groups [6, 7, 11, 12, 13]. Under the condition of character divisibility, abelian groups containing Davis-Jedwab difference sets [1, 8, 18] have also been completely characterized. As for McFarland difference sets, which are difference sets discovered by McFarland [16] with parameters

$$\begin{aligned} v &= q^n\left(\frac{q^n-1}{q-1}+1\right), \\ k &= q^{n-1}\frac{q^n-1}{q-1}, \\ \lambda &= q^{n-1}\frac{q^{n-1}-1}{q-1} \end{aligned}$$

for some positive integer n and some prime power q, the characterizations of abelian groups containing these difference sets with character divisibility property are complete only for q odd [14, 15] or $q = 4$ [8, 14, 15]. This paper is a contribution towards the understanding of abelian groups containing McFarland difference sets with q even and $q \geqslant 8$. Specifically we study McFarland difference sets with $q \geqslant 8$ a power of 2 and $n = 2$. The main result of this paper is

Theorem 1.1. *There exist* $(2^{2m+1}(2^{m-1}+1), 2^m(2^m+1), 2^m)$*-difference sets in an abelian group* G *of order* $2^{2m+1}(2^{m-1}+1)$ *with* $m \geqslant 3$ *if and only if* G *contains an elementary abelian subgroup of order* 2^{2m}.

The rest of the paper is organized as follows. In Section 2, we review the necessary background materials that are extensively used throughout this paper. These include characters of abelian groups, Frattini subgroups, group algebras and building sets introduced by Davis and Jedwab [8]. The connection between building sets and the difference sets we are interested in will also be established. In Section 3 we study subsets of abelian groups with certain character sum properties. In Section 4 we apply results obtained in Section 3 to study transversals of subgroups with certain character sum properties. We introduce two types of transversals that are important in constructing building sets for the McFarland difference sets. Section 5 is devoted to the proof of Theorem 1.1.

We thank Jim Davis for constructive suggestions and comments.

2. Preliminaries

In this section we review some of the important background materials used in this paper including characters of abelian groups, Frattini subgroups, group algebras and building sets.

2.1. **Characters.** A character χ of an finite abelian group G is a homomorphism from G to the multiplicative group of the non-zero complex numbers $\mathbb{C}^*$. The set of all characters of G is a group under the point-wise multiplication. This group is called the dual of G and is denoted by $\widehat{G}$. The identity of $\widehat{G}$, the character χ of G such that $\chi(g) = 1$ for every $g \in G$, is called the principal character of G and is denoted by $\hat{1}_G$. The group $\widehat{G}$ is isomorphic to G. If H is a subgroup of G, then restriction of a character of G to H is a character of H and this restriction map

$$\text{Res} : \widehat{G} \to \widehat{H}$$

is a group homomorphism. The kernel Ker(Res) of the restriction map consists of those characters χ of G which are principal on H, i.e. $H \leqslant \text{Ker}(\chi)$. Hence

$$\text{Ker}(\text{Res}) \cong \widehat{G/H}.$$

This implies that the restriction map Res is surjective as G is finite. The consequence of this surjectivity is that every character χ of H can be extended to a character of G and there are exactly G/H different ways of extending χ from H to G. The extendibility of characters of H to G will be frequently used in this paper. If $G = H_1 \times H_2$, then every character of G is the product of a character of H_1 and a character of H_2, i.e. $\widehat{G} = \widehat{H_1} \times \widehat{H_2}$. To give a flavor of how we use these facts in the subsequent sections, we prove the following simple lemma.

Lemma 2.1. *For any two elements $1 \neq g_1 \in G$ and $1 \neq g_2 \in G$, there is a character $\chi \in \widehat{G}$ such that $\chi(g_1) \neq 1$ and $\chi(g_2) \neq 1$.*

Proof. Let $K = \langle g_1 \rangle \cap \langle g_2 \rangle$. If $K \neq \{1\}$, then extension of any non-principal characters of K to G gives the required characters of G. If $K = \{1\}$, consider $H = \langle g_1, g_2 \rangle \cong \langle g_1 \rangle \times \langle g_2 \rangle$. Take a non-principal character of $\langle g_1 \rangle$ and a non-principal character of $\langle g_2 \rangle$, their product is a character of H and can be extended to G to obtain the required character. $\square$

2.2. **Frattini subgroup.** An element g of a group G is said to be a non-generator if $G = \langle X \rangle$ whenever $G = \langle g, X \rangle$ for any subset X of G. The set of all non-generators of a group G is a subgroup. This subgroup is called Frattini subgroup and is denoted by $\Phi(G)$. The Frattini subgroup $\Phi(G)$ of G coincides with the intersection of all proper maximum subgroups of G. If G is a finite p-group, then $\Phi(G)$ is the smallest normal subgroup N of G such that G/N is an elementary abelian p-group. If G is an abelian p-group, then obviously $\Phi(G)$ is the homomorphic image of the p-th power map of G. The groups G we will be investigating are 2-groups of exponent at most 4. The Frattini subgroup of such a group G is $\Phi(G) = \{g^2 \mid g \in G\}$ and it is elementary abelian. The characters of G that are in the Frattini subgroup of the dual of G are precisely those which are principal on the maximum elementary abelian subgroup of G, i.e. $\Phi(\widehat{G}) = \{\chi \in \widehat{G} \mid \chi(x) = 1 \text{ for all } x \in G \text{ with } x^2 = 1\}$.

2.3. **Group algebras of abelian groups.** Let G be an abelian group. The group algebra $\mathbb{C}[G]$ of the group G with complex number coefficients is the set of formal sums

$$\sum_{g \in G} a_g g,$$

where $a_g \in \mathbb{C}$ is the coefficient of g in the formal sum, endowed with the addition

$$(\sum_{g\in G} a_g g) + (\sum_{g\in G} b_g g) = \sum_{g\in G}(a_g + b_g)g,$$

and multiplication

$$(\sum_{g\in G} a_g g)(\sum_{g\in G} b_g g) = \sum_{g\in G}(\sum_{h\in G} a_h b_{gh^{-1}})g.$$

It is clear that $\mathbb{C}[G]$ is a $\mathbb{C}$-algebra. For any subset X of G we often identify X with the group algebra element

$$X = \sum_{g\in X} g$$

and define $X^{(t)}$ to be

$$X^{(t)} = \sum_{g\in X} g^t$$

for any integer $t \in \mathbb{Z}$. For example, if G is an abelian 2-group, then $G^2 = |G|G$ and $G^{(2)} = |G/\Phi(G)|\Phi(G)$ in $\mathbb{C}[G]$. A k-element subset D of G is a (v, k, λ) difference set if and only if

$$DD^{(-1)} = (k - \lambda) + \lambda G$$

in $\mathbb{C}[G]$. If $f : G_1 \to G_2$ is a group homomorphism, then f induces an algebra homomorphism $f : \mathbb{C}[G_1] \to \mathbb{C}[G_2]$ by $\mathbb{C}$-linearly extending f from G_1 to $\mathbb{C}[G_1]$. Also if $G \cong H_1 \times H_2$ and $\chi \in \widehat{H_2}$, then χ also induces an $\mathbb{C}$-algebra homomorphism $\chi : \mathbb{C}[G] \to \mathbb{C}[H_1]$. When H_1 is trivial, every character χ of G induces a $\mathbb{C}$-algebra homomorphism $\chi : \mathbb{C}[G] \to \mathbb{C}$. By the Fourier inversion formula, two elements X and Y in $\mathbb{C}[G]$ are equal if and only if $\chi(X) = \chi(Y)$ in $\mathbb{C}$ for all $\chi \in \widehat{G}$. In particular, one has

Lemma 2.2. *A k-element subset D of an abelian group G of order v is a (v, k, λ)-difference set if and only if for every non-principal $\chi \in \widehat{G}$, $|\chi(D)| = \sqrt{k - \lambda}$.*

More generally, if $G \cong H_1 \times H_2$ and X and Y are in $\mathbb{C}[G]$, then $X = Y$ if and only if $\chi(X) = \chi(Y)$ in $\mathbb{C}[H_1]$ for all $\chi \in \widehat{H_2}$.

2.4. **Building sets.** In [8], Davis and Jedwab introduced the notion of "building sets". This clever idea unifies the constructions of McFarland [16], Spence [19] and many others on Hadamard difference sets, and led to the discovery of new difference sets in [1, 4, 5, 8]. Given an abelian group G and a subgroup N of G, A family of subsets E_1, E_2, $\cdots$, E_m of G is called (k, n, m)-building sets relative to N if

(i) $|E_i| = k$, $i = 1, 2, \cdots, m$,
(ii) for very character χ of G which is trivial on N but non-trivial on G, $\chi(E_i) = 0$ for all i, and
(iii) for very character χ of G which is non-trivial on N, there is exactly one i with $|\chi(E_i)| = n$ and $\chi(E_j) = 0$ for all $j \neq i$.

For an abelian group G of order $2^{2m+1}(2^{m-1}+1)$, the prime 2 is self conjugate. By Mann Test, if D is a $(2^{2m+1}(2^{m-1}+1), 2^m(2^m+1), 2^m)$-difference set contained in G, then 2^m divides $\chi(D)$ for every $\chi \in \widehat{G}$, i.e. D has character divisibility property. From the results of [14, 15], The Sylow 2-subgroup of G is of exponent at most 4. The following lemma shows that the existence of $(2^{2m+1}(2^{m-1}+1), 2^m(2^m+1), 2^m)$-difference sets in an abelian group G of order $2^{2m+1}(2^{m-1}+1)$ is equivalent to that of $(2^{m+1}, 2^m, 2^{m-1})$-building sets in the sylow 2-subgroup of G relative to some subgroup of order 2^m.

Lemma 2.3. *Let G be an abelian of order $2^{2m+1}(2^{m-1}+1)$ whose Sylow 2-subgroup has exponent no larger than 4. The group G contains $(2^{2m+1}(2^{m-1}+1), 2^m(2^m+1), 2^m)$-difference sets if and only if its Sylow 2-subgroup contains $(2^{m+1}, 2^m, 2^{m-1})$-building sets relative to some subgroup N of order 2^m.*

Proof. Let K be the Sylow 2-subgroup of G and H the subgroup of G of order $2^{m-1}+1$. Then $G \cong K \times H$. Suppose D is a $(2^{2m+1}(2^{m-1}+1), 2^m(2^m+1), 2^m)$-difference sets in G. The set D can be written as

$$D = E_0 + h_1E_1 + h_2E_2 + \cdots + h_{2^{m-1}}E_{2^{m-1}},$$

where $H = \{1, h_1, h_2, \cdots, h_{2^{m-1}}\}$ and E_0, E_1, $\cdots$, $E_{2^{m-1}}$ are subsets of K. For every $\chi \in \widehat{K}$ and every $0 \leqslant i \leqslant 2^{m-1}$, by the Fourier inversion formular, one has

$$\chi(E_i) \quad = \quad \frac{1}{2^{m-1}+1}\sum_{\phi\in\widehat{H}}(\chi\phi)(D)\phi(h_i^{-1}).$$

Since $2^m|(\chi\phi)(D)$ and 2^m is relatively prime to $2^{m-1}+1$, one gets $2^m|\chi(E_i)$ for all $\chi \in \widehat{K}$ and all $0 \leqslant i \leqslant 2^{m-1}$. Also when χ is principal on K, the size

$$\begin{aligned}
|E_i| &= \frac{1}{2^{m-1}+1}\sum_{\phi\in\widehat{H}}(\hat{1}_K\phi)(D)\phi(h_i^{-1}) \\
&\leqslant \frac{1}{2^{m-1}+1}\sum_{\phi\in\widehat{H}}|(\hat{1}_K\phi)(D)\phi(h_i^{-1})| \\
&= \frac{1}{2^{m-1}+1}(2^m(2^m+1) + 2^{m-1}2^m) \\
&= 2^m(3 - \frac{2}{2^{m-1}+1})
\end{aligned}$$

for all $0 \leqslant i \leqslant 2^{m-1}$. Since $2^m||E_i| = \hat{1}_K(E_i)$, one has $|E_i| \leqslant 2^{m+1}$ for all $0 \leqslant i \leqslant 2^{m-1}$. The fact that $|D| = 2^m(2^m+1)$ implies that $|E_i| = 2^{m+1}$ for all i except one, say $i = 0$, with $|E_0| = 2^m$. When χ is non-principal on K, the absolute value

$$\begin{aligned}
|\chi(E_i)| &= \frac{1}{2^{m-1}+1}|\sum_{\phi\in\widehat{H}}(\chi\phi)(D)\phi(h_i^{-1})| \\
&\leqslant \frac{1}{2^{m-1}+1}\sum_{\phi\in\widehat{H}}|(\chi\phi)(D)\phi(h_i^{-1})| \\
&= 2^m
\end{aligned}$$

for all $0 \leqslant i \leqslant 2^{m-1}$. Since $2^m|\chi(E_i)$ and $\chi(E_i) \in \mathbb{Z}[\sqrt{-1}]$ as K is of exponent no larger than 4, one has $|\chi(E_i)| = 2^m$ or 0 for all non-principal $\chi \in \widehat{K}$ and all $0 \leqslant i \leqslant 2^{m-1}$. This implies that E_0 is a coset of a subgroup, say N, of order 2^m in K. From $DD^{(-1)} = 2^{2m} + 2^mG$, one gets

$$\sum_{i=0}^{2^{m-1}} E_iE_i^{(-1)} \quad = \quad 2^{2m} + 2^mK$$

and

$$\sum_{i=0}^{2^{m-1}} |\chi(E_i)|^2 = \sum_{i=0}^{2^{m-1}} \chi(E_i E_i^{(-1)}) = 2^{2m}$$

for every non-principal $\chi \in \widehat{K}$. Hence E_1, E_2, $\cdots$, $E_{2^{m-1}}$ form $(2^{m+1}, 2^m, 2^{m-1})$-building sets in K relative to N. Conversely, if K contains $(2^{m+1}, 2^m, 2^{m-1})$-building sets E_1, E_2, $\cdots$, $E_{2^{m-1}}$ relative to N of order 2^m, then it is easy to check that when $E_0 = N$ and

$$D = E_0 + h_1 E_1 + h_2 E_2 + \cdots + h_{2^{m-1}} E_{2^{m-1}},$$

one has $|\chi(D)| = 2^m$ for all non-principal $\chi \in \widehat{G}$ and therefore, by Lemma 2.2, D is a $(2^{2m+1}(2^{m-1}+1), 2^m(2^m+1), 2^m)$-difference set in G. □

3. SUBSETS IN ABELIAN 2-GROUPS WITH CERTAIN CHARACTER SUM PROPERTIES

In this section we study subsets E in abelian 2-groups G of exponent no larger than 4 such that $|\chi(E)| = |E|/2$ or 0 for all non-principal $\chi \in \widehat{G}$. The results in this section will used in subsequent sections in investigating $(2^{m+1}, 2^m, 2^{m-1})$-building sets. We start with the following two simple technical lemmas. The first one shows that $E \cap K$ inherits character divisibility from E while the second demonstrates that E contains a coset of a subgroup of order $|E|/4$ if $|\chi(E)| = |E|/2$ for some character χ of order 2 in $\widehat{G}$.

Lemma 3.1. *Let G be a finite abelian group of even order and E a subset of G. Suppose E has the property that there is a positive integer l such that for every character $\chi \in \widehat{G}$, the character sum $\chi(E) \equiv 0 \pmod{2l}$. Then for every index 2 subgroup K of G, the subset $E \cap K$ has the property that for every $\chi \in \widehat{K}$, the character sum $\chi(E \cap K) \equiv 0 \pmod{l}$.*

Proof. Let K be an index 2 subgroup of G and $x \in G \setminus K$. The set E can be decomposed into

$$E = (E \cap K) + x(x^{-1}E \cap K).$$

For every character $\chi \in \widehat{K}$, there are two characters χ^+ and χ^- in $\widehat{G}$ such that $\chi^+|_K = \chi^-|_K = \chi$ and $\chi^+(x) + \chi^-(x) = 0$. Applying χ^+ and χ^- to E, one gets

$$\chi^+(E) = \chi(E \cap K) + \chi^+(x)\chi(x^{-1}E \cap K) \equiv 0 \pmod{2l},$$
$$\chi^-(E) = \chi(E \cap K) + \chi^-(x)\chi(x^{-1}E \cap K) \equiv 0 \pmod{2l}.$$

The sum of the two equations ensures that $\chi(E \cap K) \equiv 0 \pmod{l}$. □

Lemma 3.2. *Let G be an abelian 2-group of exponent no larger than 4 and E a subset of G such that $|E| \equiv 0 \pmod 4$. If the subset E satisfies*

1. *for every character $\chi \in \widehat{G}$, the value $\chi(E) \equiv 0 \pmod{|E|/2}$, and*
2. *there is a character $\phi \in \widehat{G}$ of order 2 such that $|\phi(E)| = |E|/2$,*

then E contains a coset of a subgroup of order $|E|/4$ in G .

Proof. Let K be the kernel of ϕ. Since ϕ is of order 2, the subgroup K is of index 2 in G. Let $x \in G \setminus K$. Then $E = (E \cap K) + x(x^{-1}E \cap K)$ and $\phi(E) = |E \cap K| - |x^{-1}E \cap K| = \pm|E|/2$. Hence either $|E \cap K| = |E|/4$ or $|x^{-1}E \cap K| = |E|/4$. By Lemma 3.1, $\chi(E \cap K) \equiv \chi(x^{-1}E \cap K) \equiv 0 \pmod{|E|/4}$ for all $\chi \in \widehat{K}$. Therefore E contains a coset of a subgroup of order $|E|/4$ in G. □

The next lemma shows that the subgroup $(\mathbb{Z}/2\mathbb{Z})^3$ in $(\mathbb{Z}/4\mathbb{Z})^3$ has no transversal E such that $\chi(E) \equiv 0 \pmod 4$ for all characters of $(\mathbb{Z}/4\mathbb{Z})^3$. This is actually a crucial fact of our proof of Theorem 1.1.

Lemma 3.3. *Let $G \cong (\mathbb{Z}/4\mathbb{Z})^3$ and N the maximum elementary abelian subgroup of G. There is no transversal E of N in G such that $\chi(E) \equiv 0 \pmod 4$ for every $\chi \in \widehat{G}$.*

Proof. Let $G = \langle x\rangle \times \langle y\rangle \times \langle z\rangle$, where x, y and z are of order 4. Up to equivalence by isomorphisms and translations of G, we may assume that

$$E = 1 + x + y + z + xyw_{xy} + yzw_{yz} + zxw_{zx} + xyzw_{xyz},$$

where w_{xy}, w_{yz}, w_{zx} and w_{xyz} are elements in N. Since $\chi(E) \equiv 0 \pmod 4$ for every $\chi \in \widehat{G}$, one can check that

$$\prod_{g\in E} \chi(g) = 1$$

for every $\chi \in \widehat{G}$. Therefore

$$\prod_{g\in E} g = w_{xy}w_{yz}w_{zx}w_{xyz} = 1$$

and $w_{xyz} = w_{xy}w_{yz}w_{zx}$. If we apply character χ such that $\chi(x) = \pm\sqrt{-1}$, $\chi(y) = \pm 1$ and $\chi(z) = \pm 1$, we find that $1 + \chi(y) + \chi(z) + \chi(y)\chi(z)\chi(w_{yz}) \equiv 0 \pmod 4$ and $1 + \chi(y)\chi(w_{xy}) + \chi(z)\chi(w_{zx}) + \chi(y)\chi(z)\chi(w_{xyz}) \equiv 0 \pmod 4$. This is equivalent to $\chi(w_{yz}) = 1$ because $a_1 + a_2 + a_3 + a_4 \equiv 0 \pmod 4$ if and only if $a_1a_2a_3a_4 = 1$ when $a_i = \pm 1$ for $i = 1, 2, 3, 4$. Therefore

$$w_{yz} \in \{1, y^2, z^2, y^2z^2\}.$$

Similarly,

$$\begin{aligned}
&w_{zx} \in \{1, x^2, z^2, x^2z^2\},\\
&w_{xy} \in \{1, x^2, y^2, x^2y^2\},\\
&w_{yz}w_{xyz} \in \{1, x^2, y^2z^2, x^2y^2z^2\},\\
&w_{zx}w_{xyz} \in \{1, y^2, z^2x^2, x^2y^2z^2\},\\
&w_{xy}w_{xyz} \in \{1, z^2, x^2y^2, x^2y^2z^2\},\\
&w_{xyz} \in \{x^2, y^2, z^2, x^2y^2z^2\}.
\end{aligned}$$

Therefore we need w_{xy}, w_{yz}, w_{xz} and w_{xyz} in N satisfy

$$w_{xy} \in \{1, x^2, y^2, x^2y^2\} \cap w_{xyz}\{1, z^2, x^2y^2, x^2y^2z^2\}, \tag{1}$$

$$w_{yz} \in \{1, y^2, z^2, y^2z^2\} \cap w_{xyz}\{1, x^2, y^2z^2, x^2y^2z^2\}, \tag{2}$$

$$w_{zx} \in \{1, x^2, z^2, x^2z^2\} \cap w_{xyz}\{1, y^2, z^2x^2, x^2y^2z^2\}, \tag{3}$$

$$w_{xyz} \in \{x^2, y^2, z^2, x^2y^2z^2\} \tag{4}$$

$$w_{xy}w_{yz}w_{xz}w_{xyz} = 1. \tag{5}$$

One can easily check that there is no w_{xyz} in $\{x^2, y^2, z^2, x^2y^2z^2\}$ that can produce w_{xy}, w_{yz} and w_{zx} that satisfy all conditions above. □

For a p-group G and a normal subgroup N of G, it is clear that $\Phi(N)$ is contained in $N \cap \Phi(G)$ as $\Phi(N)$ is the smallest normal subgroup X of N such that N/X is elementary abelian while $N/(N \cap \Phi(G))$ is a subgroup of $G/\Phi(G)$ which is elementary abelian. An interesting consequence of Lemma 3.3, which we prove as the following corollary, is that if a subgroup N of an abelian 2-group G of exponent no larger than 4 has a transversal E with the property that $\chi(E) \equiv 0 \pmod{|E|/2}$ for all $\chi \in \widehat{G}$, then $\Phi(N)$ is at most index 4 in $N \cap \Phi(G)$. that is

Corollary 3.4. *Let G be an abelian 2-group of exponent no larger than 4 and N a subgroup of G. Let E be a transversal of N in G. If for every character $\chi \in \widehat{G}$, the value $\chi(E) \equiv 0 \pmod{|E|/2}$, then $|(N \cap \Phi(G))/\Phi(N)| \leqslant 4$.*

Proof. If $|(N \cap \Phi(G))/\Phi(N)| \geqslant 8$, there is a subgroup $K \cong (\mathbb{Z}/4\mathbb{Z})^3$ of G such that $N \cap K \cong (\mathbb{Z}/2\mathbb{Z})^3$ and $\Phi(N) \cap K = \{1\}$. Then the group N has a subgroup N' such that $N' \cap K = \{1\}$ and $N = (N \cap K)N'$. In the quotient group G/N', the image $\overline{E}$ of E is a transversal of the image $\overline{N} = N/N' \cong N \cap K < \overline{K} \cong K$. The transversal also has the property that for every character $\chi \in \widehat{\overline{G}}$, the value $\chi(\overline{E}) \equiv 0 \pmod{|\overline{E}|/2}$ since E has. Now we can find a sequence of subgroups $\overline{K} = H_0 < H_1 < H_2 \cdots < H_{k-1} < H_k = \overline{G}$, where k is the positive integer satisfies $|\overline{G}| = 2^{k+6}$, such that $|H_i/H_{i-1}| = 2$ for $i = 1, 2, \cdots, k$. Applying Lemma 3.1 successively to the subgroups $H_{k-1}, H_{k-2}, \cdots, H_0 = \overline{K}$, one finds that $\overline{E} \cap \overline{K}$ has the property that $\chi(\overline{E} \cap \overline{K}) \equiv 0 \pmod{4}$ for every character $\chi \in \widehat{\overline{K}}$. Also $\overline{E} \cap \overline{K}$ is a transversal of $\overline{N}$ in $\overline{K}$. This contradicts Lemma 3.3. $\square$

As an application of Corollary 3.4, we prove the following lemma which shows that every set in $(2^{m+1}, 2^m, 2^{m-1})$-building sets satisfies the conditions of Lemma 3.2.

Lemma 3.5. *Let $m \geqslant 3$ be an integer. Let G be an abelian group of order 2^{2m+1} and of exponent at most 4, and E a subset of G of size 2^{m+1}. If for every non-principal character $\chi \in \widehat{G}$, $|\chi(E)| = 2^m$ or 0, then there is a character χ of order 2 in $\widehat{G}$ such that $|\chi(E)| = |E|/2$.*

Proof. Suppose to the contrary. Then $\chi(E) = 0$ for every character χ of order 2 in $\widehat{G}$, i.e. every $\chi \in \widehat{G}$ with $\Phi(G) \leqslant \mathrm{Ker}(\chi)$. Let $\overline{G} = G/\Phi(G)$. Then the image $\overline{E}$ of E in $\overline{G}$ must be a multiple of the whole group. Hence $|\overline{G}| = |G/\Phi(G)| \leqslant |E| = 2^{m+1}$, or equivalently, $|\Phi(G)| \geqslant |G|/|E| = 2^m$. Since G is of exponent at most 4, $|\Phi(G)| \leqslant 2^m$. Therefore $|\Phi(G)| = 2^m$ and E is a transversal of $\Phi(G)$ in G, which contradicts Corollary 3.4. $\square$

4. Transversals in abelian 2-groups with certain character sum properties

In this section we introduce two types of transversals and show that they arise frequently in the building sets that we are interested in. Let G be an abelian 2-group of exponent no larger than 4 and N a subgroup of G. Let E be a transversal of N in G. The two types of transversals that we want to investigate are

(I) There are two subgroups H_1 and H_2 of G such that $|H_1| = |H_2| = |E|/2$ and $E = aH_1 + bH_2$ for some a and b in G;

(II) There is a subgroup H of G of order $|E|/8$ and an 8-element subset E' of G such that $E = HE'$.

The next two two lemmas give descriptions of the structure of type (I) transversals. The first one is valid without any restrictions while the second is for transversal E with the property that $|\chi(E)| = |E|/2$ or 0.

Lemma 4.1. *If E is of type* (I)*, then $|H_1 \cap N| = |H_2 \cap N| = 1$, $H_1N = H_2N$ and $a^{-1}b \notin H_1N$.*

Proof. Since E is a transversal of N in G, it is obvious that $|H_1 \cap N| = |H_2 \cap N| = 1$. Since $a^{-1}E$ is also a transversal of N in G, the intersection $NH_1 \cap a^{-1}E = H_1$ is a transversal of N in NH_1, and therefore $a^{-1}bH_2 \cap NH_1 = \emptyset$. Hence $a^{-1}b \notin NH_1$. Since NH_1 is an index 2 subgroup of $G = NE$, one finds that the subgroup $H_2 \subseteq ab^{-1}(G \setminus NH_1) = NH_1$. Since $|H_1N| = |H_2N|$, one gets $H_1N = H_2N$. □

Lemma 4.2. *Let E be a transversal of type* (I). *If E has the property that $|\chi(E)| = |E|/2$ or 0 for every non-principal character $\chi \in \widehat{G}$, then $N \leqslant \langle H_1, H_2\rangle$, $|H_1 \cap H_2| = |E|/(2|N|)$ and $H_1/(H_1 \cap H_2) \cong H_2/(H_1 \cap H_2) \cong N$.*

Proof. Let $H = \langle H_1, H_2\rangle$, by Lemma 4.1, $H \leqslant NH_1 = NH_2$. From the absolute values of character sums of $a^{-1}E$, we know that $G = \langle a^{-1}E\rangle = \langle a^{-1}b, H\rangle$. We show that $N \leqslant H$. If there is an $x \in N$ such that $x \notin H$, then $x = (a^{-1}b)^s h$ for some integer $1 \leqslant s \leqslant 3$ and some $h \in H$. By Lemma 4.1, $s \neq 1$ or 3 as $(a^{-1}b)^s = xh^{-1} \in NH = NH_1 = NH_2$. This implies that $s = 2$, $a^{-1}b$ is an element of order 4 and $|\langle a^{-1}b\rangle \cap H| = 1$. Hence there is a non-principal character $\chi \in \widehat{G}$ with $\chi((a^{-1}b)^2) = -1$ and χ is principal on H, so that $|\chi(E)| = \sqrt{2}|E|/2$. This contradicts the assumption that $|\chi(E)| = |E|/2$ or 0 for every non-principal character $\chi \in \widehat{G}$. Therefore $N \leqslant H$ and $H = NH_1 = NH_2$. ¿From $|\langle H_1, H_2\rangle| = |H_1||H_2|/|H_1 \cap H_2|$, one gets $|H_1 \cap H_2| = |E|/(2|N|)$. Now consider the group $\overline{G} = G/(H_1 \cap H_2)$ and the image $\overline{E}$ of E in $\overline{G}$. Clearly, $\overline{E} = |H_1 \cap H_2|(\overline{a}\overline{H}_1 + \overline{b}\overline{H}_2)$ and $\overline{a}\overline{H}_1 + \overline{b}\overline{H}_2$ is a transversal of $\overline{N}$ in $\overline{G}$ which has the property that for every non-principal character $\chi \in \widehat{\overline{G}}$, the absolute value $|\chi(\overline{a}\overline{H}_1 + \overline{b}\overline{H}_2)| = |\overline{a}\overline{H}_1 + \overline{b}\overline{H}_2|/2$ or 0, where $\overline{H}_1 = H_1/(H_1 \cap H_2)$, $\overline{H}_2 = H_2/(H_1 \cap H_2)$ and $\overline{N} \cong N$ is the image of N in $\overline{G}$. Since $|\overline{H}_1 \cap \overline{H}_2| = |\overline{H}_1 \cap \overline{N}| = |\overline{H}_2 \cap \overline{N}| = 1$ and $\overline{H}_1\overline{H}_2 = \overline{H}_1\overline{N} = \overline{H}_2\overline{N}$, one gets $\overline{H}_1 \cong \overline{H}_2 \cong \overline{N} \cong N$. □

The next two lemmas show that for a type (II) transversal E of N in G, there is a restriction on the size of the subgroup N when we impose the condition $\chi(E) = |E|/2$ or 0 for every non-principal $\chi \in \widehat{G}$.

Lemma 4.3. *Let G be a finite abelian 2-group and N a subgroup of G. If a transversal E of N in G satisfies*

(1) *$|E| \leqslant 8$, and*

(2) *$|\chi(E)| = |E|/2$ or 0 for every non-principal character $\chi \in \widehat{G}$,*

then $|N| \leqslant |E|$.

Proof. Consider the group ring element

$$X = E^2E^{(-1)} - \frac{|E|^2}{4}E \in \mathbb{Z}[G].$$

Since $\chi(X) = 0$ for every non-principal character $\chi \in \widehat{G}$, $X = \lambda G$ for some integer λ. Applying the principal character of G, one gets $\lambda = (3|E|^3)/(4|G|) = (3|E|^2)/(4|N|)$ and $|N|$ divides $|E|^2/4$ as G is a 2-group. When $|E| = 2$ or 4, we clearly have

$|N| \leqslant |E|$. When $|E| = 8$, if λ is even, then $|N|$ divides $|E|^2/8$ and therefore $|N| \leqslant |E|$. If λ is odd, then $\lambda = 3$ and

$$E^2 E^{(-1)} = \frac{|E|^2}{4} E + 3G.$$

¿From $E^2 = E^{(2)} + 2U$ for some $U \in \mathbb{Z}[G]$, one gets $E^{(2)}E^{(-1)} - G = 2V$ for some $V \in \mathbb{Z}[G]$. This implies that for every $g \in G$, the coefficient of g in $E^{(2)}E^{(-1)}$ is odd, and therefore $|G| \leqslant |E^{(2)}E^{(-1)}| = |E|^2$ and $|N| \leqslant |E|$. □

Corollary 4.4. *Let E be of type* (II). *If E has the property that $|\chi(E)| = |E|/2$ or 0 for every non-principal character $\chi \in \widehat{G}$, then $|N| \leqslant 8$.*

Proof. Since E is a transversal of N in G and also it is the union of 8 cosets of the subgroup H, one must have $|H \cap N| = 1$ and E' is a transversal of NH in G. Let $\overline{E}$ be the image of E in $\overline{G} = G/H$. Then $\overline{E} = |H|\overline{E}'$, where $\overline{E}'$ is the image of E' in $\overline{G}$. Since E' is a transversal of NH in G, its image $\overline{E}'$ contains 8 elements and is a transversal of $\overline{N} \cong N$ in $\overline{G}$. Because E has the property that $|\chi(E)| = |E|/2$ or 0 for every non-principal character $\chi \in \widehat{G}$, the transversal $\overline{E}'$ also has the property that $|\chi(\overline{E}')| = |\overline{E}'|/2$ or 0 for every non-principal character $\chi \in \widehat{\overline{G}}$. By Lemma 4.3, $|N| \leqslant 8$. □

If the set E in Lemma 3.2 is a transversal of a subgroup N of G, then by a translation, we can assume that such a transversal contains a subgroup A of G with $|A| = |E|/4$. Obviously $|A \cap N| = 1$ as A is contained in a transversal of N, and $[G : NA] = 4$. The following two lemmas show that E is of type (I) or (II) when $G/NA \cong (\mathbb{Z}/2\mathbb{Z})^2$ and is of type (I) when $G/NA \cong \mathbb{Z}/4\mathbb{Z}$ and N is elementary.

Lemma 4.5. *Let G be an abelian 2-group of exponent no larger than 4 and N a subgroup of G such that $|N| \geqslant 8$. Let E be a transversal of N in G which contains a subgroup A of G such that $|A| = |E|/4$. If*

(1) *for every character $\chi \in \widehat{G}$, the absolute value $|\chi(E)| = |E|/2$ or 0, and*
(2) *$G/NA \cong (\mathbb{Z}/2\mathbb{Z})^2$,*

then E must be of type (I) *or* (II).

Proof. Let $\{1, x, y, xy\}$ be a transversal of NA in G. Since $G/NA \cong (\mathbb{Z}/2\mathbb{Z})^2$, we have three index 2 subgroups $H_1 = (1 + x)NA$, $H_2 = (1 + y)NA$ and $H_3 = (1+xy)NA$ of G. By Lemma 3.1, the set $E \cap H_i$ has the property that $\chi(E \cap H_i) \equiv 0 \pmod{|E|/4}$ for all i. Note that E is a transversal of N in G and $N < H_i$ for all i, therefore $E \cap H_i$ is a transversal of N in H_i and $|E \cap H_i| = |E|/2$ for all i. Since $E \cap NA = A$, one gets

$$|E \cap xNA| = |E \cap yNA| = |E \cap xyNA| = |E|/4$$

and

$$\chi(E \cap xNA) \equiv \chi(E \cap yNA) \equiv \chi(E \cap xyNA) \equiv 0 \pmod{|E|/4}.$$

This implies that $E \cap xNA$, $E \cap yNA$ and $E \cap xyNA$ are cosets of subgroups of order $|E|/4$ in G. Hence we can assume that $E \cap xNA = xuA_1$, $E \cap yNA = yvA_2$ and $E \cap xyNA = xywA_3$, where A_i is a subgroup of NA such that $|A_i| = |E|/4$ and $|A_i \cap N| = 1$ for $i = 1$, 2 and 3, and u, v and w are in N as $NA_i = NA$ for all i, and $E = A + xuA_1 + yvA_2 + xywA_3$.

If two of the four groups A, A_1, A_2 and A_3 are the same, we claim that E is of type (I). Without loss of generality, we assume that $A = A_2$. Then $A_1 = A_3$ because $\chi(E) \equiv 0 (\mathrm{mod}\ |E|/2)$ implies that χ is principal on even number of the groups A, A_1, A_2 and A_3 for every $\chi \in \widehat{NA}$. Therefor $A + yuA_2 = A(1 + yu)$ and $xvA_1 + xywA_3 = xvA_1(1 + ywv^{-1})$. We now want to show that $(yu)^2 \in A$ or $(ywv^{-1})^2 \in A_1$, and it follows that $A(1+yu)$ is a group or $A_1(1+ywv^{-1})$ is a group, and subsequently it implies that both of them are groups as $\chi(E) \equiv 0 (\mathrm{mod}\ |E|/2)$ for all $\chi \in \widehat{G}$, and hence E is of type (I). Suppose to the contrary. Then $(yu)^2 \notin A$ and $(ywv^{-1})^2 \notin A_1$. Let's look at $(yu)^2(ywv^{-1})^2$. Since $(yu)^2(ywv^{-1})^2 = (uwv^{-1})^2 \in N$ and $A \cap N = \{1\}$, if $(yu)^2(ywv^{-1})^2 \neq 1$, then we can find a $\chi \in \widehat{G}$ such that $\chi((yu)^2(ywv^{-1})^2) = \chi((yu)^2) = -1$ and χ is principal on A, which implies that

$$\chi(A(1 + yu)) = (1 \pm \sqrt{-1})|E|/4 \text{ and } \chi(xvA_1(1 + ywv^{-1})) \equiv 0 \pmod{|E|/2}$$

as $\chi((ywv^{-1})^2) = 1$. This contradicts the condition that $\chi(E) \equiv 0 \pmod{|E|/2}$. So $(yu)^2(ywv^{-1})^2 = 1$ and $(yu)^2 = (ywv^{-1})^2$. Now $(yu)^2 = (ywv^{-1})^2$ can not be in $\langle a, A_1\rangle$ because if $(yu)^2 = (ywv^{-1})^2 \in \langle a, A_1\rangle$, then there is a $\chi \in \widehat{G}$ such that $\chi((yu)^2) = -1$ and χ is principal on A as $(yu)^2 \notin A$, which would ensure $\chi(A_1) = 0$ as $(yu)^2 = (ywv^{-1})^2 \in \langle a, A_1\rangle$ and $\chi(E) = (1\pm\sqrt{-1})|E|/4$, contradicting $\chi(E) \equiv 0 \pmod{|E|/2}$. Now since $(yu)^2 = (ywv^{-1})^2 \notin \langle a, A_1\rangle$, one must have $A = A_1$ because, again, if $A \neq A_1$, then there is a $\chi \in \widehat{G}$ such that $\chi((yu)^2) = -1$, χ is principal on A but not principal on A_1 as $A \neq A_1$, which implies that $\chi(E) = (1 \pm \sqrt{-1})|E|/4$ contradicting $\chi(E) \equiv 0 \pmod{|E|/2}$. Hence $E = A(1+xu+yv+xyw)$. In the group $\overline{G} = G/A$, the image of $\{1, xu, yv, xyw\}$ form a transversal of $\overline{N} \cong N$ in $\overline{G}$ and has the property that $|\chi(1+\overline{xu}+\overline{yv}+\overline{xyw})| = 2$ or 0 for all non-principal character $\chi \in \widehat{\overline{G}}$. By Lemma 4.3, $N \leqslant 4$, contradicting $N \geqslant 8$.

If no two of A, A_1, A_2 and A_3 are the same, let $A_0 = A$, we claim that

$$|\bigcap_{i=0}^{3} A_i| = |E|/8.$$

Let $H = \bigcap_{i=0}^{3} A_i$. By taking the quotient G/H, we may assume that $H = \{1\}$ and we need to show that $|E| = 8$. From $|H| = 1$, we can see that the intersections of any three of the four groups A, A_1, A_2 and A_3 are trivial. This is because if there are three groups, say A_1, A_2 and A_3 such that $\bigcap_{i=1}^{3} A_i \neq \{1\}$, then there is a character $\chi \in \widehat{G}$ such that χ is non-principal on $\bigcap_{i=1}^{3} A_i$ and principal on $A_0 = A$ as $A \cap (\bigcap_{i=1}^{3} A_i) = \{1\}$, which implies $|\chi(E)| = |E|/4$, a contradiction. We now calculate the sizes of pairwise intersections of the 4 groups. Let $F = NA_0 = NA_1 = NA_2 = NA_3$ and consider the images $\overline{A}_1$, $\overline{A}_2$ and $\overline{A}_3$ of A_1, A_2 and A_3 respectively in the group $F/A_0 \cong N$. None of these three images is trivial as $A_i \nleqslant A_0$ for all $i \neq 0$. Also every character $\chi \in \widehat{F/A_0}$ must be trivial on at least one of the three images as such a χ of F/A_0 can be extended to a character of G which is principal on A_0. If there are two images, say $\overline{A}_1$ and $\overline{A}_2$, such that $\overline{A}_1 \cap \overline{A}_2 \neq \{1\}$, then for any $1 \neq u \in \overline{A}_1 \cap \overline{A}_2$ and $1 \neq v \in \overline{A}_3$, by Lemma 2.1, there is a character $\chi \in \widehat{F/A_0}$ such that $\chi(u) \neq 1$ and $\chi(v) \neq 1$, which implies there are characters which are non-principal on all three images and this is not possible. Hence the pairwise intersections of the three images must be trivial. Let $1 \neq a_i \in \overline{A_i}$ for $i = 1, 2$ and 3. Since there is no character χ such that $\chi(a_i) \neq 1$ for all i, the set $\{1, a_1, a_2, a_3\}$ must be a group isomorphic to $(\mathbb{Z}/2\mathbb{Z})^2$. Fix a_1 and a_2, we can see that $\overline{A}_3 = \{1, a_3\}$. Similarly, $\overline{A}_i = \{1, a_i\}$ for all $i = 1$, 2 and 3. Hence $|A_i/(A_0 \cap A_i)| = |\overline{A}_i| = 2$ for all i, and similarly, $|A_i/(A_i \cap A_j)| = 2$ for all $i \neq j$. Since $|(A_i \cap A_j) \cap (A_i \cap A_k)| = 1$ when i, j and k are distinct, either $A_i \cong (\mathbb{Z}/2\mathbb{Z})^2$ for all i, or $A_i \cong \mathbb{Z}/2\mathbb{Z}$ for all i as these are the only groups containing index 2 subgroups with trivial intersections. If $A_i \cong (\mathbb{Z}/2\mathbb{Z})^2$, then $\langle a_0, A_1, A_2, A_3\rangle \cong (\mathbb{Z}/2\mathbb{Z})^3$, which is impossible as there are characters that are trivial on exactly one of these A_i's since they are hyperplanes of $\langle a_0, A_1, A_2, A_3\rangle$. So $|A_i| = 2$ for $i = 0$, 1, 2 and 3 and $|E| = 8$. □

Lemma 4.6. *Let G be an abelian 2-group of exponent no larger than 4 and N an elementary abelian subgroup of G such that $|N| \geqslant 8$. Let E be a transversal of N in G which contains a subgroup A of G such that $|A| = |E|/4$. If*

1. *for every character $\chi \in \widehat{G}$, the absolute value $|\chi(E)| = |E|/2$ or 0, and*
2. *$G/NA \cong \mathbb{Z}/4\mathbb{Z}$,*

then E must be of type (I).

Proof. If $G/NA \cong \mathbb{Z}/4\mathbb{Z}$, we may further assume that $E = A + xA_1 + x^2A_2 + x^3A_3 = (A + x^2A_2) + x(A_1 + x^2A_3)$, where A_1, A_2 and A_3 are transversals of N in NA and x is an element of order 4 in G such that $|\langle x\rangle \cap (NA)| = 1$. Since $A + x^2A_2 = E \cap \langle x^2\rangle NA$, By Lemma 3.1, $\chi(A + x^2A_2) \equiv 0 \pmod{|E|/4}$ for all $\chi \in \widehat{\langle x^2\rangle NA}$. It follows that $A_2 = uB$ for some subgroup B of order $|E|/4$ in NA with $|B \cap N| = 1$ and $NA = NB$, and $u \in N$ because A is a group of order $|E|/4$. We now show that $A = B$. Let $F = NA = NB$. If $A \neq B$, then there is a $\chi \in \widehat{G}$ such that $\chi(A) \neq \chi(B)$ and $\chi(x^2) = -1$ as $|\langle x\rangle \cap F| = 1$. Since $F/A \cong F/B \cong N$ is elementary abelian and either A or B is contained in the kernel of χ, the character $\chi|_F$ must be of order 2. Therefore $\chi(A_1)$ and $\chi(A_3)$ are real, and consequently, $\chi(A_1 + x^2A_3)$ is real, which contradicts $|\chi(E)| = |E|/2$ or 0 for all non-principal $\chi \in \widehat{G}$. Hence $A = B$ and $A + x^2A_2 = A(1 + ux^2)$ is a group as ux^2 is of order 2. Now for every $\chi \in \widehat{G}$,

$$\chi(x(A_1 + x^2A_3)) = \chi(E) - \chi(A + x^2A_2) \equiv 0 \pmod{|E|/2}.$$

This implies that $A_1 + x^2A_3$ is a coset of a subgroup of order $|E|/2$ in G and E is of type (I). □

We now change the conditions imposed on G/NA to conditions on characters and obtained following two corollaries which are actually the ones used in the proof of the main theorem.

Corollary 4.7. *Let G be an abelian 2-group of exponent no larger than 4 and N an elementary abelian subgroup of G such that $|N| \geqslant 8$. If a transversal E of N in G satisfies*

1. *for every non-principal character $\chi \in \widehat{G}$, the absolute value $|\chi(E)| = |E|/2$ or 0, and*
2. *there is a character $\phi \in \widehat{G}$ of order 2 such that $|\phi(E)| = |E|/2$,*

then E must be of type (I) *or* (II).

Proof. By Lemma 3.2, we may assume that the set E contains a subgroup A of order $|E|/4$ in G. Since E is a transversal of N in G, $|A \cap N| = 1$ and $[G : NA] = 4$. By Lemma 4.5 and 4.6, the transversal E is either of type (I) or (II). □

Corollary 4.8. *Let G be an abelian 2-group of exponent 4 and N a subgroup of G such that $|N| \geqslant 8$ and the exponent of N is 4. If a transversal E of N in G satisfies*

1. *for every non-principal character $\chi \in \widehat{G}$, the absolute value $|\chi(E)| = |E|/2$ or 0, and*
2. *there is a character $\phi \in \Phi(\widehat{G})$ such that $|\phi(E)| = |E|/2$,*

then E must be of type (I) *or* (II).

Proof. Let $K = \mathrm{Ker}(\phi)$ be the kernel of ϕ. Since $\phi \in \Phi(\widehat{G})$, ϕ is of order 2 and $|G/K| = 2$. By the proof of Lemma 3.2, we can assume that $E \cap K$ is a subgroup of K of order $|E|/4$. We know that $G = K + xK$ for some $x \notin K$. the element x is of order 4 as $\phi \in \Phi(\widehat{G})$ and we can also assume that $x \in N$ as E is a transversal of N in G. We now show that there is an index 2 subgroup H of K such that $|\langle x^2 \rangle \cap H| = 1$. This is equivalent to show that $x^2 \notin \Phi(K)$, i.e. there is no $y \in K$ such that $y^2 = x^2$. Suppose that there is an $y \in K$ such that $y^2 = x^2$. Then $(xy)^2 = 1$. This implies that $xy \in K$ as $\phi \in \Phi(\widehat{G})$, and hence $x \in K$, a contradiction. Let H be a subgroup of K such that $|\langle x^2 \rangle \cap H| = 1$ and $K = \langle x^2 \rangle H$. The group $E \cap K$ is not contained in H because if $E \cap K \leqslant H$, then there is a character $\chi \in \widehat{G}$ such that χ is principal on H and $\chi(x^2) = -1$, which implies that

$$\chi(E) = \chi(E \cap K) + \chi(x)\chi(x^{-1}E \cap K) = (|E|/4) \pm \chi(x^{-1}E \cap K)\sqrt{-1}$$

and $\chi(x^{-1}E \cap K)$ is real as $\chi|_K$ is of order 2 since $H \leqslant \mathrm{Ker}(\chi)$, contradicting the condition that $|\chi(E)| = |E|/2$ or 0 for every non-principal character $\chi \in \widehat{G}$. Hence $E \cap H$ is an index 2 subgroup of $E \cap K$ and $E \cap K = (E \cap H)(1 + x^2 y)$ for some $y \in H$ such that $y \notin E$ but $y^2 \in E$. Let $F = (E \cap H)(1 + y)$. F is a subgroup of H of order $|E|/4$. Let H_1 be an index 2 subgroup of H such that $E \cap H \leqslant H_1$. If $y \notin H_1$, then there is a $\chi \in \widehat{G}$ such that χ is principal on H_1 but not on H and $\chi(x^2) = -1$. Such a χ is real on K as $K = \langle x^2 \rangle H$ and $\chi|_H$ is real. Hence, again

$$\chi(E) = \chi(E \cap K) + \chi(x)\chi(x^{-1}E \cap K) = (|E|/4) \pm \chi(x^{-1}E \cap K)\sqrt{-1}$$

contradicting the condition that $|\chi(E)| = |E|/2$ or 0 for every non-principal character $\chi \in \widehat{G}$. Therefore $y \in H_1$ for every index 2 subgroup H_1 of H that contains $E \cap H$. This is equivalent to that the rank of $H/(E \cap H)$ is equal to that of $H/(E \cap H)(1 + y)$, or the image of y in $H/(E \cap H)$ is in $\Phi(H/(E \cap H))$, i.e. $y = st^2$ for some $s \in E \cap H$ and $t \in H$ but $|(E \cap H) \cap \langle t \rangle| = 1$. Therefore $F = (E \cap H)(1 + y) = (E \cap H)(1 + t^2)$ and $E \cap K = (E \cap H)(1 + x^2 t^2)$. Now $t^2 \notin N(E \cap H)$ because if $t^2 = uv$ with $u \in N$ and $v \in E \cap H$, then

$$E \cap K = (E \cap H)(1 + x^2 t^2) = (E \cap H)(1 + x^2 uv) = (E \cap H)(1 + x^2 u),$$

which is impossible as E is a transversal of N in G and $x^2 u \in N$. Hence $|\langle t \rangle \cap N(E \cap H)| = 1$ and $G/N(E \cap H) \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$. Since $N(E \cap K) = N(E \cap H)(1 + x^2 t^2) = N(E \cap H)(1 + t^2)$ as $x^2 \in N$, one gets $G/N(E \cap K) \cong (\mathbb{Z}/2\mathbb{Z})^2$ and, by Lemma 4.5, E is of type (I) or (II). □

We finish this section with a lemma that shows the groups involved in the type (I) transversals of $(2^{m+1}, 2^m, 2^{m-1})$-building sets form some kind of orthogonal system.

Lemma 4.9. *Let G be an abelian group of order 2^{2m+1} and of exponent at most 4 and N a subgroup of G of order 2^m. Let $E_1 = a_1 H_1 + a_2 H_2$, $E_2 = a_3 H_3 + a_4 H_4$, $\cdots$,*

$E_k = a_{2k-1}H_{2k-1} + a_{2k}H_{2k}$ be a family of type (I) *transversals of N in G, where H_i are subgroups of order 2^m in G and $H_i \cap N = \{1\}$ for all $i = 1, 2, \cdots, k$. If for every non-principal character $\chi \in \widehat{G}$, there is at most one E_i such that $|\chi(E_i)| = 2^m$, and $\chi(E_j) = 0$ for all $j \neq i$, then $H_s \cap H_t = \{1\}$ and $NH_s = NH_t = H_sH_t$ for all $1 \leqslant s \neq t \leqslant 2k$.*

Proof. According to Lemma 4.2 and the condition that for every non-principal character $\chi \in \widehat{G}$, there is at most one E_i such that $|\chi(E_i)| = 2^m$, and $\chi(E_j) = 0$ for all $j \neq i$, we can see that χ can be principal on at most one H_i, $1 \leqslant i \leqslant k$ when χ is non-principal on N. If χ is non-principal on N and principal on H_s for some $1 \leqslant s \leqslant k$, then χ is a character of $\overline{G} = G/H_s$. The image $\overline{N}$ of N in $\overline{G}$ is isomorphic to N as $H_s \cap N = \{1\}$. For every $t \neq s$, the group $\overline{N}$ must be contained in the image group $\overline{H}_t$ of H_t because otherwise there is a $\chi \in \widehat{\overline{G}}$ such that χ is principal on $\overline{H}_t$ and non-principal on $\overline{N}$. Therefore $\overline{H}_t = \overline{N}$ and $H_tH_s = NH_s$ for all $1 \leqslant s \neq t \leqslant 2k$. □

5. PROOF OF THEOREM 1.1

In this we prove Theorem 1.1. We start with two simple lemmas and their proofs, which we omit, are easy exercises of applications of characters.

Lemma 5.1. *If an abelian group G of order $2^{m+1}n$ contains $(2^{m+1}, 2^m, 2^{m-1})$-building sets relative to a subgroup N of order n in G, then for any subgroup H of N, the group G/H contains $(2^{m+1}, 2^m, 2^{m-1})$-building sets relative to a subgroup N/H.*

Lemma 5.2. *Let G be a group of order 2^{m+n+1} and N a subgroup of G of order 2^n. If $E_1, E_2, E_3, \cdots, E_{2^{m-1}}$ form $(2^{m+1}, 2^m, 2^{m-1})$-building sets in G relative to N, then*

$$\sum_{k=1}^{2^{m-1}} E_k E_k^{(-1)} = 2^{2m} + 2^{2m-n}(G - N).$$

The next lemma excludes type (II) transversals from any $(2^{m+1}, 2^m, 2^{m-1})$-building sets in a group of order 2^{2m+1}.

Lemma 5.3. *Let G be a group of order 2^{2m+1} and $m \geqslant 3$. There are no type* (II) *transversals in any $(2^{m+1}, 2^m, 2^{m-1})$-building sets relative to any subgroup of order 2^m in G.*

Proof. If $E_1, E_2, \cdots, E_{2^{m-1}}$ form $(2^{m+1}, 2^m, 2^{m-1})$-building sets relative to a subgroup of order 2^m in G and, say, E_1 is of type (II), then $E_1 = HE'$ for some non-trivial subgroup H of G and some subset E' of G with $|E'| = 8$. Let $x \in H$ and $x \neq 1$. Then the coefficient of x in $E_1E_1^{(-1)}$ is $|E_1| = 2^{m+1}$, which contradicts Lemma 5.2. □

We now prove Theorem 1.1.

Proof of Theorem 1.1. The existence part of the theorem is given in [10, 16]. By Lemma 2.3 and results in [14, 15], we need to prove that any abelian 2-group G of order 2^{2m+1} and of exponent no larger than 4 containing $(2^{m+1}, 2^m, 2^{m-1})$-building sets relative to a subgroup N of order 2^m must contain an elementary abelian subgroup of index 2 in G. Let E_1, E_2, $\cdots$, $E_{2^{m-1}}$ be $(2^{m+1}, 2^m, 2^{m-1})$-building sets relative to N in G. We are going to show that N is an elementary abelian subgroup of G.

Suppose N is not elementary abelian. Then there are characters $\chi \in \Phi(\widehat{G})$ which are not principal on N. By the definition of building sets, E_1, E_2, $\cdots$, $E_{2^{m-1}}$ are transversals of N in G. By Corollary 4.8 and Lemma 5.3, there are type (I) transversals among E_1, E_2, $\cdots$, $E_{2^{m-1}}$. Assume that $E_1 = a_1H_1 + a_2H_2$, $E_2 = a_3H_3 + a_4H_4$, $\cdots$, $E_k = a_{2k-1}H_{2k-1} + a_{2k}H_{2k}$, $k \leqslant 2^{m-1}$ are of type (I) transversals and the rest are not.

Claim 1: $k = |\Phi(N)|/2$.

Let $H = NH_1$. By Lemma 4.9, $H = NH_s = H_sH_t$ for all $1 \leqslant s \neq t \leqslant 2k$ and H is an index 2 subgroup of G. Let $H_0 = N$. Since $H_i < H$ and $H \cong H_i \times H_i$ for all $0 \leqslant i \leqslant 2k$, $\Phi(H_i) < \Phi(H)$ and $\Phi(H) \cong \Phi(H_i) \times \Phi(H_i)$ for all $0 \leqslant i \leqslant 2k$. Since $\Phi(H_i) \cong \Phi(N)$ as $H_i \cong N$ and $(\Phi(H_i)\setminus\{1\}) \cap (\Phi(H_j)\setminus\{1\}) = \emptyset$ as $H_i \cap H_j = \{1\}$ for all $0 \leqslant i \neq j \leqslant 2k$, there can be at most $|\Phi(H)\setminus\{1\}|/|\Phi(N)\setminus\{1\}|$ such subgroups of H_i, i.e. $2k+1 \leqslant (|\Phi(H)|-1)/(|\Phi(N)|-1) = (|\Phi(N)|^2-1)/(|\Phi(N)|-1) = |\Phi(N)|+1$. Therefore $k \leqslant |\Phi(N)|/2$. On the other hand, for each type (I) transversal E_i, a $\chi \in \Phi(\widehat{G})$ has $|\chi(E_i)| = 2^m$ if and only if χ is non-principal on N, principal on the maximum elementary abelian subgroup of G and principal on exactly one of H_{2i-1} and H_{2i}. There are exactly $2[(|\Phi(G)|/|\Phi(N)|) - (|\Phi(G)|/|\Phi(N)|^2)]$ such $\chi \in \Phi(\widehat{G})$. The group $\Phi(\widehat{G})$ contains $|\Phi(G)| - (|\Phi(G)|/|\Phi(N)|)$ characters which are not principal on N. Therefore, by By Corollary 4.8 and Lemma 5.3, there should be at least

$$\frac{|\Phi(G)| - (|\Phi(G)|/|\Phi(N)|)}{2[(|\Phi(G)|/|\Phi(N)|) - (|\Phi(G)|/|\Phi(N)|^2)]} = \frac{|\Phi(N)|}{2}$$

E_i's of type (I). Hence $k = |\Phi(N)|/2$.

Claim 2: $k \leqslant 2$**.**

Since $k = |\Phi(N)|/2$, the sets $\Phi(H_i) \setminus \{1\}$, $i = 0, 1, \cdots, |\Phi(N)|$ form a partition of $\Phi(H)\setminus\{1\}$. By Lemma 3.5, for each $|\Phi(N)|/2 \leqslant i \leqslant |N|/2$, E_i contains a coset of a subgroup, say K_i, of order 2^{m-1} of G. By Lemma 5.2, $K_i \cap \Phi(H) = \{1\}$. Therefore $K_i \cap H$ is an elementary abelian group. Since $K_i \cap N = \{1\}$ as E_i is a transversal of N in G, one has

$$\mathrm{Rank}(H) \geqslant \mathrm{Rank}(K_i \cap H) + \mathrm{Rank}(N) = \mathrm{Rank}(K_i \cap H) + [\mathrm{Rank}(H)/2].$$

Therefore $\mathrm{Rank}(H) \geqslant 2\mathrm{Rank}(K_i \cap H)$. Since H is of index 2 in G and $|K_i| = 2^{m-1}$, the order $|K_i \cap H| \geqslant 2^{m-2}$ and the rank $\mathrm{Rank}(K_i \cap H) \geqslant m-2$. This implies $\mathrm{Rank}(H) \geqslant 2(m-2) = 2m-4$ and $\mathrm{Rank}(\Phi(H)) \leqslant 4$ as $|H| = 2^{2m}$. Hence $k = |\Phi(N)|/2 = \sqrt{|\Phi(H)|}/2 \leqslant 2$.

Claim 3: $k = 1$**.**

If $k = 2$, then $N \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^{m-4}$, $H \cong (\mathbb{Z}/4\mathbb{Z})^4 \times (\mathbb{Z}/2\mathbb{Z})^{2m-8}$. Also for each $i > 2$, we have $K_i \cap H \cong (\mathbb{Z}/2\mathbb{Z})^{m-2}$ and $N(K_i \cap H) \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^{2m-6}$. Let $x \in K_i$ such that $x \notin H$. Then

$$G/NK_i = \langle H, x\rangle / \langle N(K_i \cap H), x\rangle \cong H/N(K_i \cap H) \cong (\mathbb{Z}/2\mathbb{Z})^2.$$

By Lemma 4.5 and 5.3, E_i is of type (I), contradicting the assumption that E_i is not of type (I) for $|\Phi(N)|/2 \leqslant i \leqslant |N|/2$. Hence $k = 1$.

Since $k = |\Phi(N)|/2 = 1$, one has

$$N \cong \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^{m-2}, \text{ and } H \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^{2m-4}.$$

Also by Lemma 4.5 and 5.3, $G/NK_i \cong \mathbb{Z}/4\mathbb{Z}$ for $i \geqslant 2$. From $|G/H| = 2$, one has either $\mathrm{Rank}(\Phi(G)) = 3$ and $G \cong (\mathbb{Z}/4\mathbb{Z})^3 \times (\mathbb{Z}/2\mathbb{Z})^{2m-5}$, or $\mathrm{Rank}(\Phi(G)) = 2$ and $G \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^{2m-3}$.

Claim 4: $\mathrm{Rank}(\Phi(G)) \neq 3$.

If $\mathrm{Rank}(\Phi(G)) = 3$ and $G \cong (\mathbb{Z}/4\mathbb{Z})^3 \times (\mathbb{Z}/2\mathbb{Z})^{2m-5}$, let's consider the group $\overline{G} = G/\Phi(N) \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^{2m-4}$. In $\overline{G}$, the image $\overline{N} = N/\Phi(N)$ of N is an elementary abelian group of order 2^{m-1}, and, by Lemma 5.1, the images $\overline{E}_1$, $\overline{E}_2$, $\cdots$, $\overline{E}_{2^{m-1}}$ of E_1, E_2, $\cdots$, $E_{2^{m-1}}$ form $(2^{m+1}, 2^m, 2^{m-1})$-building sets relative to $\overline{N}$ in $\overline{G}$. The transversal $\overline{E}_1$ is of type (I) since E_1 is. For $i \geqslant 2$, $\overline{E}_i$ contains the group $\overline{K}_i$, the image of K_i in G, and $\overline{K}_i \cong K_i$ as $K_i \cap N = \{1\}$. Since $\overline{G}/\overline{NK}_i \cong G/NK_i \cong \mathbb{Z}/4\mathbb{Z}$ and $\overline{N}$ is elementary abelian, by Lemma 4.6, the transversal $\overline{E}_i$ is of type (I) for every $i \geqslant 2$. We can now assume that $\overline{E}_i = {a'}_{2i-1}{H'}_{2i-1} + {a'}_{2i}{H'}_{2i}$ for some subgroups ${H'}_{2i-1}$ and ${H'}_{2i}$ of order 2^m in $\overline{G}$ and some ${a'}_{2i-1}$ and ${a'}_{2i}$ in $\overline{G}$ for $i = 1, 2, \cdots, 2^{m-1}$. The group ${H'}_j \cap \overline{N} = \{1\}$ for all j as $\overline{E}_i$ is a transversal of $\overline{N}$. Hence ${H'}_j$ is not elementary abelian, i.e. $\Phi({H'}_j)$ is non-trivial, for all j because $\mathrm{Rank}(\overline{G}) = 2m - 2$ while $|{H'}_j| = 2^m$, $|\overline{N}| = 2^{m-1}$ and $\overline{N}$ is elementary abelian. Also by Lemma 4.2, ${H'}_{2i-1} \cap {H'}_{2i}$ is of order 2 and contains $\Phi({H'}_{2i-1})$ and $\Phi({H'}_{2i})$, and therefore ${H'}_{2i-1} \cap {H'}_{2i} = \Phi({H'}_{2i-1}) = \Phi({H'}_{2i})$ is contained in $\Phi(\overline{G})$ for all i. By Lemma 5.2, $({H'}_{2s-1} \cap {H'}_{2s}) \neq ({H'}_{2t-1} \cap {H'}_{2t})$ for all $s \neq t$, i.e. ${H'}_{2i-1} \cap {H'}_{2i}$ are distinct subgroups of order 2 in $\Phi(\overline{G})$. Since $|\Phi(\overline{G})| = 4$, the group $\Phi(\overline{G})$ only has three subgroups of order 2. Hence $2^{m-1} \leqslant 3$ and $m \leqslant 2$, contradicting $m \geqslant 3$.

Claim 4: $\mathrm{Rank}(\Phi(G)) \neq 2$.

If $\mathrm{Rank}(\Phi(G)) = 2$ and $G \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^{2m-3}$, let N' be an elementary abelian subgroup of N of order 2^{m-2} such that $N' \cap \Phi(N) = \{1\}$. By Lemma 5.1, the images $\overline{E}_1$, $\overline{E}_2$, $\cdots$, $\overline{E}_{2^{m-1}}$ of E_1, E_2, $\cdots$, $E_{2^{m-1}}$ in $\overline{G} = G/N'$ form $(2^{m+1}, 2^m, 2^{m-1})$-building sets relative to $\overline{N} = N/N' \cong \mathbb{Z}/4\mathbb{Z}$. Moreover the transversal $\overline{E}_1$ of $\overline{N}$ in $\overline{G}$ is of type (I) since E_1 is and $\overline{E}_1 = \overline{a}_1\overline{H}_1 + \overline{a}_2\overline{H}_2$, where $\overline{a}_1$, $\overline{a}_2$, $\overline{H}_1$ and $\overline{H}_2$ are images of a_1, a_2, H_1 and H_2 respectively in $\overline{G}$. We now show that intersection $\overline{H}_1 \cap \overline{H}_2$ is an elementary abelian group of order 2^{m-2}. Since $H = H_1H_2 = NH_1 = NH_2$ and $N \cap H_1 = N \cap H_2 = H_1 \cap H_2 = \{1\}$, there are isomorphisms $f_1 : N \to H_1$ and $f_2 : N \to H_2$ such that $N = \{f_1(x)f_2(x) \in H_1H_2 = H \mid x \in N\}$. Hence

$$\begin{aligned} N'H_1 &= \{h_1 f_2(y) \in H_1H_2 = H \mid h_1 \in H_1, y \in N'\}, \\ N'H_2 &= \{f_1(x) h_2 \in H_1H_2 = H \mid x \in N', h_2 \in H_2\} \end{aligned}$$

and $N'H_1 \cap N'H_2 = \{f_1(x)f_2(y) \in H_1H_2 = H \mid x, y \in N'\} \cong N' \times N'$. This implies that

$$\overline{H}_1 \cap \overline{H}_2 = (N'H_1 \cap N'H_2)/N' \cong N'$$

is an elementary abelian group of order 2^{m-2}. Let $F = \overline{H}_1 \cap \overline{H}_2$ and $\overline{N} = \langle z \rangle$. By Lemma 4.2, we can assume that there is an element x of order 4 in $\overline{G}$ such that $\overline{H}_1 = \langle F, x \rangle$ and $\overline{H}_2 = \langle F, xz \rangle$. The group $\overline{H} = \langle \overline{H}_1, \overline{H}_2 \rangle = \langle F, x, z \rangle$ is an index 2 subgroup of $\overline{G}$. Let $y \in \overline{G} \backslash \overline{H}$. Then $y^2 = 1$. Let $K = \langle F, x, y \rangle$. Then $K \cap \overline{N} = \{1\}$. For each $2 \leqslant i \leqslant 2^{m-1}$, the set $\overline{E}_i$ must have the form

$$\overline{E}_i = F_{i0} + F_{i1}z + F_{i2}z^2 + F_{i3}z^3,$$

where F_{i0}, F_{i1}, F_{i2} and F_{i3} are disjoint subsets of K such that $F_{i0}+F_{i1}+F_{i2}+F_{i3} = K$. Since for any character $\chi \in \widehat{\overline{G}}$ with $\chi(z) = -1$, $|\chi(\overline{E}_1)| = 2^m$ if and only if χ is principal on $\langle F, x^2\rangle$, one finds that $F_{20} + F_{22}$, $F_{30} + F_{32}$, $\cdots$, $F_{2^{m-1}0} + F_{2^{m-1}2}$ form $(2^m$, 2^{m-1}, $2^{m-1} - 1)$-building sets in K relative to $\langle F, x^2\rangle$, and hence we can further assume that

$$F_{i0} + F_{i2} = A_{i0} + A_{i2}x$$

for some subsets A_{i0} and A_{i2} of sizes 2^{m-1} in $\langle y, F, x^2\rangle$ for all $i = 2, 3, \cdots, 2^{m-1}$. Clearly, $F_{i1} + F_{i3}$ is the complement of $F_{i0} + F_{i2}$ in K, therefore $F_{i1} + F_{i3} = A_{i1} + A_{i3}x$, where $A_{i1} = \langle y, F, x^2\rangle - A_{i0}$ and $A_{i3} = \langle y, F, x^2\rangle - A_{i2}$ for all $i \geqslant 2$. Let

$$\overline{E}_i = B_{i0} + B_{i1}z + B_{i2}x + B_{i3}xz,$$

where

$$\begin{aligned} B_{i0} &= \overline{E}_i \cap \langle y, F, x^2, z^2\rangle, \\ B_{i1} &= z^{-1}\overline{E}_i \cap \langle y, F, x^2, z^2\rangle, \\ B_{i2} &= x^{-1}\overline{E}_i \cap \langle y, F, x^2, z^2\rangle, \\ B_{i3} &= (xz)^{-1}\overline{E}_i \cap \langle y, F, x^2, z^2\rangle. \end{aligned}$$

If $p : \langle y, F, x^2, z^2\rangle \to \langle y, F, x^2\rangle$ is the natural projection, then $p(B_{is}) = A_{is}$ for all $i \geqslant 2$ and $s = 0, 1, 2, 3$, i.e. B_{i0}, B_{i1}, B_{i2} and B_{i3} are the lift-backs of A_{i0}, A_{i1}, A_{i2} and A_{i3} from $\langle y, F, x^2\rangle$ to $\langle y, F, x^2, z^2\rangle$. Clearly, $B_{i0} + B_{i2}x = F_{i0} + F_{i2}z^2$ and $B_{i1} + B_{i3}x = F_{i1} + F_{i3}z^2$ for all $i \geqslant 2$. Now we want to evaluate the coefficients of x^2z^2 in $\overline{E}_i\overline{E}_i^{(-1)}$ for $i \geqslant 2$. If the coefficient of x^2z^2 in $\overline{E}_i\overline{E}_i^{(-1)}$ is not 0 for some $i \geqslant 2$, then $\overline{E}_i$ contains a coset of $\langle x^2z^2\rangle$. Obviously, such a coset must be in one of B_{is}, for some $0 \leqslant s \leqslant 3$. Without loss of generality, we can assume that B_{i0} contains a coset of $\langle x^2z^2\rangle$. Let $\chi \in \widehat{\langle z^2\rangle}$ such that $\chi(z^2) = -1$. Then χ induces a $\mathbb{C}$-algebra homomorphism, which we denoted again by χ, $\chi : \mathbb{C}[K \times \langle z^2\rangle] \to \mathbb{C}[K]$. Let $\rho : \mathbb{C}[K] \to \mathbb{C}[K/\langle x^2\rangle]$ be the $\mathbb{C}$-algebra homomorphism induced by natural projection $K \to K/\langle x^2\rangle$. Then $\rho\chi(F_{i0} + F_{i2}z^2) \in \mathbb{C}[K/\langle x^2\rangle]$. For every character $\phi \in \widehat{K/\langle x^2\rangle}$, $\phi\rho\chi$ is a character of $K \times \langle z^2\rangle$ with $\phi\rho\chi(z^2) = -1$ and $\phi\rho\chi(x^2) = 1$. It can be extended to $\overline{G}$ and

$$\phi\rho\chi(\overline{E}_i) = \phi\rho\chi(F_{i0} + F_{i2}z^2) \pm \phi\rho\chi(F_{i1} + F_{i3}z^2)\sqrt{-1}.$$

Since $\phi\rho\chi(F_{i0} + F_{i2}z^2)$ and $\phi\rho\chi(F_{i1} + F_{i3}z^2)$ are real as $\phi\rho\chi(x^2) = 1$, one has $|\phi\rho\chi(F_{i0} + F_{i2}z^2)| = 2^m$ or 0. On the other hand, $F_{i0} + F_{i2}z^2$ contains cosets of $\langle x^2z^2\rangle$ and they are in the kernel of $\rho\chi$. Therefore $|\phi\rho\chi(F_{i0} + F_{i2}z^2)| \leqslant |F_{i0} + F_{i2}z^2| - 2(\text{number of cosets of } \langle x^2z^2\rangle) < 2^m$. Hence $\phi(\rho\chi(F_{i0} + F_{i2}z^2)) = 0$ for all $\phi \in \widehat{K/\langle x^2\rangle}$. This implies that $\rho\chi(F_{i0} + F_{i2}z^2) = 0$ in $\mathbb{C}[K/\langle x^2\rangle]$, or equivalently, $\rho(F_{i0}) = \rho(F_{i2})$. The fact that F_{i0} and F_{i2} are disjoint subsets of K implies that $F_{i0} = x^2F_{i2}$ and $F_{i0}+F_{i2}z^2 = (1+x^2z^2)F_{i0}$. Since $B_{i0}+B_{i2}x = F_{i0}+F_{i2}z^2$, the sets B_{i0} and B_{i2} are unions of 2^{m-2} cosets of $\langle x^2z^2\rangle$ and A_{i0} and A_{i2} are unions of 2^{m-2} cosets of $\langle x^2\rangle$ as A_{is} is the image of B_{is} under the natural projection $\overline{G} \to \overline{G}/\langle z^2\rangle$. Since $A_{i1} = \langle y, F, x^2\rangle - A_{i0}$ and $A_{i3} = \langle y, F, x^2\rangle - A_{i2}$, A_{i1} and A_{i3} are also unions of 2^{m-2} cosets of $\langle x^2\rangle$. Clearly these cosets in A_{i1} and A_{i3} must come from cosets of $\langle x^2\rangle$ in B_{i1} and B_{i3} or cosets of $\langle x^2z^2\rangle$ in B_{i1} and B_{i3}. Therefore B_{i1} and B_{i3} are in unions of cosets of $\langle x^2\rangle$ and cosets of $\langle x^2z^2\rangle$. If B_{i1} and B_{i3} contains no cosets of $\langle x^2z^2\rangle$, i.e. $B_{i1} + B_{i3}x = F_{i1} + F_{i3}z^2$ is the union of 2^{m-1} cosets of $\langle x^2\rangle$, let

$\chi_1 \in \widehat{\langle x^2 \rangle}$ such that $\chi_1(x^2) = -1$ and $\chi_2 \in \widehat{\langle z^2 \rangle}$ such that $\chi_2(z^2) = -1$ and $\chi_1\chi_2$ and χ_2 induces $\mathbb{C}$-algebra homomorphisms

$$\chi_1\chi_2 : \mathbb{C}[\langle y, F, x^2, z^2 \rangle] \to \mathbb{C}[\langle y, F \rangle]$$

and

$$\chi_2 : \mathbb{C}[\langle K, z^2 \rangle] \to \mathbb{C}[K].$$

For every character $\phi \in \widehat{\langle y, F \rangle}$, one has $\phi\chi_1\chi_2 \in \widehat{\langle y, F, x^2, z^2 \rangle}$ which can be extended to $\overline{G}$. Applying such an extension to $\overline{E}_i$, one finds that $\phi\chi_1\chi_2(B_{i0}) = \phi\chi_1\chi_2(B_{i2}) = 0$ for all ϕ. Hence $\chi_1\chi_2(B_{i0}) = \chi_1\chi_2(B_{i2}) = 0$ in $\mathbb{C}[\langle y, F \rangle]$. On the other hand, for $\chi_0 \in \widehat{\langle x^2 \rangle}$ with $\chi_0(x^2) = 1$, one also has $\chi_0\chi_2(B_{i0}) = \chi_0\chi_2(B_{i2}) = 0$ as B_{i0} and B_{i2} are unions of 2^{m-2} cosets of $\langle x^2z^2 \rangle$. Therefore $\chi_2(B_{i0}) = \chi_2(B_{i2}) = 0$ in $\mathbb{C}[\langle y, F, x^2 \rangle]$. This implies that $\chi_2(F_{i0} + F_{i2}z^2) = 0$ in $\mathbb{C}[K]$ or $F_{i0} = F_{i2}$ in K. Since F_{i0} and F_{i2} are disjoint, $F_{i0} = F_{i2} = \emptyset$, which contradicts the assumption that B_{i0} contains a coset of $\langle x^2z^2 \rangle$. Hence B_{i1} or B_{i3} contains cosets of $\langle x^2z^2 \rangle$, i.e. $F_{i1} + F_{i3}z^2$ contains cosets of $\langle x^2z^2 \rangle$. Using arguments similar to those for $F_{i0} + F_{i2}z^2$, one can show that $F_{i1} + F_{i3}z^2$ is the union of 2^{m-1} cosets of $\langle x^2z^2 \rangle$. Therefore $\overline{E}_i$ is the union of 2^m cosets of $\langle x^2z^2 \rangle$, i.e. the coefficient of x^2z^2 in $\overline{E}_i\overline{E}_i^{(-1)}$ is $|\overline{E}_i| = 2^{m+1}$. Thus for each $i \geqslant 2$, the coefficient of x^2z^2 in $\overline{E}_i\overline{E}_i^{(-1)}$ is either 0 or 2^{m+1}. Since the coefficient of x^2z^2 in $\overline{E}_1\overline{E}_1^{(-1)}$ is 2^m, the coefficient of x^2z^2 in

$$\sum_{k=1}^{2^{m-1}} E_kE_k^{(-1)}$$

is congruent to $2^m \pmod{2^{m+1}}$. By Lemma 5.2, one gets $2^{2m-2} \equiv 2^m \pmod{2^{m+1}}$, a contradiction as $m \geqslant 3$.

Therefor the group N must be elementary abelian. By Lemma 3.5, Corollary 4.7, Lemma 5.3 and Lemma 4.2, the group G contains an index 2 elementary abelian subgroup $H \cong N \times N$. □

As we can see from the proof of Theorem 1.1, the construction of abelian $(2^{2m+1}(2^{m-1}+1), 2^m(2^m+1), 2^m)$-difference sets for $m \geqslant 3$ is essentially unique and is given in [10] and [16].

REFERENCES

[1] K. T. Arasu and Y. Q. Chen, A difference set in $(\mathbb{Z}/4\mathbb{Z})^3 \times \mathbb{Z}/5\mathbb{Z}$, *Des. Codes Cryptogr.* **23** (2001), 317–323.

[2] K. T. Arasu and S. K. Sehgal, Some new difference sets, *J. Combin. Theory Ser. A* **69** (1995), 170–172.

[3] T. Beth, D. Jungnickel and H. Lenz, *Design Theory*, volume 1. Cambridge University Press, Cambridge, 2nd edition, 1999.

[4] Y. Q. Chen, On the existence of abelian Hadamard difference sets and a new family of difference sets, *Finite Fields Appl.* **3** (1997), 234–256.

[5] Y. Q. Chen, A construction of difference sets, *Des. Codes Cryptogr.* **13** (1998), 247–250.

[6] J. A. Davis, A result on Dillon's conjecture in difference sets, *J. Combin. Theory Ser. A* **57** (1991), 238–242.

[7] J. A. Davis, Difference sets in abelian 2-groups, *J. Combin. Theory Ser. A* **57** (1991), 262–286.

[8] J. A. Davis and J. Jedwab, A unifying construction for difference sets, *J. Combin. Theory Ser. A* **80** (1997), 13–78.

[9] J. A. Davis and J. Jedwab, A new family of relative difference sets in 2-groups, *Des. Codes Cryptogr.* **17** (1999), 305–312.

[10] J. F. Dillon, Variations on a scheme of McFarland for noncyclic difference sets, *J. Combin. Theory Ser. A* **40** (1985), 9–21.

[11] J. F. Dillon, Difference sets in 2-groups, In: Finite Geometry and Combinatorial Designs, *Contemporary Mathematics*, **111**, (Ed. E. S. Kraemer), BirKhäuser, Boston, (1990), 65–72.

[12] J. F. Dillon, A survey of difference in 2-groups, Presented at the Marshall Hall Memorial Conference, Burlington, VT, (1990).

[13] R. G. Kraemer, Proof of a conjecture on Hadamard 2-groups, *J. Combin. Theory Ser. A* **63** (1993), 1–10.

[14] S. L. Ma and B. Schmidt, The structure of abelian groups containing McFarland difference sets, *J. Combin. Theory Ser. A* **70** (1995), 313–322.

[15] S. L. Ma and B. Schmidt, A sharp exponent Bound for McFarland difference sets with $p = 2$, *J. Combin. Theory Ser. A* **80** (1997), 347–352.

[16] R. L. McFarland, A family of difference sets in non-cyclic abelian groups, *J. Combin. Theory Ser. A* **15** (1973), 1–10.

[17] A. Pott, *Finite Geometry and Character Theory*, Vol. 1601 of Lecture Notes in Mathematics, Springer-Verlag, Berlin, Heidelberg, 1995.

[18] B. Schmidt, Nonexistence results on Chen and Davis-Jedwab difference sets, *J. Algebra* **202** (1998), 404–413.

[19] T. Spence, A family of difference sets in non-cyclic groups, *J. Combin. Theory Ser. A* **22** (1977), 103–106.

Department of Mathematics and Statistics, Wright State University, Dayton, Ohio 45435, USA

Institute for Algebra and Geometry, Otto-von-Guericke-University Magdeburg, 39016 Magdeburg, Germany